Carbohydrate Research, 224 (1992) c9–c11
Elsevier Science Publishers B.V., Amsterdam

CONTENTS

(Abstracts/contents lists published in: Chem. Abstr.; Curr. Contents: Phys., Chem. & Earth Sci.; Life Sci. Full texts are incorporated in CJELSEVIER, *a file in the Chemical Journals Online database which is available on STN® International.)*

Roles of potassium ions, acetyl and L-glyceryl groups in native gellan double helix: an X-ray study
R. Chandrasekaran, A. Radha, and V. G. Thailambal (West Lafayette, IN, U.S.A.) . . . 1

E.s.r. study of the alkaline degradation of disaccharides in methyl sulfoxide
P. Calle, A. Sanchez, and C. Sieiro (Madrid, Spain) 19

4-Thiocellooligosaccharides. Their synthesis and use as ligands for the separation of cellobiohydrolases of *Trichoderma reesei* by affinity chromatography
C. Orgeret, E. Seillier, C. Gautier, J. Defaye, and H. Driguez (Grenoble, France) 29

Novel types of acyclic *C*-nucleoside analogues *via* the reaction of hydrogen bromide in acetic acid with L-*threo*-(glycerol-1-yl)pyrazolinediones
H. A. Hamid, N. Rashed, Y. El Kilany, and E. S. H. El Ashry (Alexandria, Egypt) 41

Synthesis, structure, and transformations of 1,2,4-phosphites of α-D-xylopyranose
E. E. Nifant'ev, M. P. Koroteev, V. A. Sychev, A. R. Bekker, L. K. Vasyanina, and N. N. Nevskii (Moscow, U.S.S.R.) 49

Spacer-modified disaccharide and pseudo-trisaccharide methyl glycosides that mimic maltotriose, as competitive inhibitors for pancreatic alpha-amylase: a demonstration of the "clustering effect"
M. Blanc-Muesser, L. Vigne, H. Driguez (Grenoble, France), J. Lehmann, J. Steck, and K. Urbahns (Freiburg, Germany) 59

Preparation of higher-carbon sugars by stereoselective osmylation of related allylic alcohols
S. Jarosz (Warsaw, Poland) 73

Synthesis of a tetrasaccharide acceptor for use in the assay of UDP-Glc*p*NAc:β-D-Gal*p*-(1→4)-β-D-Glc*p*NAc (GlcNAc to Gal) β(1→3)-*N*-acetylglucosaminyltransferase activity and the pentasaccharide product that would be formed by its enzymic glycosylation
G. Srivastava and O. Hindsgaul (Edmonton, Alta., Canada) 83

Synthesis and conformation of dihistamine derivatives of cyclomaltoheptaose (β-cyclodextrin)
V. Cucinotta, F. D'Alessandro, G. Impellizzeri, and G. Vecchio (Catania, Italy) 95

Total stereo- and enantio-selective synthesis of 2,3-dideoxy-3-*C*-methylene-D-*glycero*-pentose and its ethyl furanosides
Yu. E. Raifeld, G. Ya. Vid, I. E. Mikerin, B. M. Arshava, and A. A. Nikitenko (Moscow, U.S.S.R.) 103

Synthesis of nigero-oligosaccharides
K. Takeo, S. Kitamura, and Y. Murata (Kyoto, Japan) 111

Synthesis of sialic acid-containing nucleotides sugars: CMP-sialic acid analogs
K. Ikeda, Y. Nagao, and K. Achiwa (Shizuoka, Japan) 123

In situ synthesis of ferrites in lignocellulosics
R. H. Marchessault, S. Ricard, and P. Rioux (Montreal, Que., Canada) 133

A synthetic study of methyl 3-deoxy-3-fluoro-α-D-glucopyranosides from methyl 2,3-anhydro-α-D-allopyranosides, and synthesis of 3′-deoxy-3′-fluorokanamycin A and 3′-chloro-3′-deoxykanamycin A
E. Umemura, T. Tsuchiya, Y. Kobayashi, and K. Tanaka (Kawasaki, Japan) 141

Inclusion complexes of some non-ionic surfactants with cyclomalto-oligosaccharides
T. Cserháti and J. Szejtli (Budapest, Hungary) 165

Synthesis of an *S*-(α-sialosyl)-(2→9)-*O*-(α-sialosyl)-(2→3′)-β-lactosylceramide
A. Hasegawa, H. Ogawa, H. Ishida, and M. Kiso (Gifu, Japan) 175

Synthesis of a position isomer of ganglioside GD_3 having an α-Neu5Ac-(2→9)-α-Neu5Ac linkage
A. Hasegawa, H. Ogawa, and M. Kiso (Gifu, Japan) 185

Application of cyclic sulfates in the synthesis of 6-deoxy-D-*manno*-heptopyranose derivatives
P. A. M. van der Klein and J. H. van Boom (Leiden, The Netherlands) 193

The distribution of carboxymethyl groups in *O*-(carboxymethyl)ated (1→3)-β-D-glucans and (1→3)-α-D-glucans
S. Ukai, I. Yoshida, A. Honda, K. Nagai, and T. Kiho (Gifu, Japan) 201

Structure of a heteroglycan isolated from the fungus *Omphalia lapidescens*
K. Saito, M. Nishijima, and T. Miyazaki (Tokyo, Japan) 209

New investigations of the structure of grape arabinogalactan–protein
L. Saulnier, J.-M. Brillouet, M. Moutounet (Montpellier, France), C. Hervé du Penhoat, and V. Michon (Paris, France) 219

Structure and antitumor activity of a branched (1→3)-β-D-glucan from the alkaline extract of *Amanita muscaria*
T. Kiho, M. Katsuragawa, K. Nagai, S. Ukai (Gifu, Japan), and M. Haga (Hokkaido, Japan) 237

Structure of the extracellular polysaccharide from slime-forming *Lactococcus lactis* subsp. *cremoris* SBT 0495
H. Nakajima, T. Hirota (Kawagoe, Japan), T. Toba, T. Itoh, and S. Adachi (Sendai, Japan) 245

The use of glycosides of 6- and 8-acylamino-4-methylumbelliferone in studies of the specificity and properties of human lysosomal glycolipid hydrolases
G. Ya. Wiederschain, I. K. Kozlova, G. S. Ilyina, M. A. Mikhaylova, and E. M. Beyer (Moscow, U.S.S.R.) 255

Notes

Crystal and molecular structure of 4,8-anhydro-3-deoxy-3-nitro-D-*lyxo*-D-*gluco*-nonitol
P. Köll (Oldenburg, Germany), J. Kopf, M. Morf, B. Zimmer (Hamburg, Germany), M. Petrušová, and L. Petruš (Bratislava, Czechoslovakia) 273

Preparation and n.m.r. assignments of cellulose mixed esters regioselectively substituted by acetyl and propanoyl groups
T. Iwata, J.-I. Azuma, K. Okamura (Kyoto, Japan), M. Muramoto (Osaka, Japan), and B. Chun (Kanagawa, Japan) 277

An n.m.r. study of the 1,3,4,6- and 1,2,3,6-tetra-acetates of β-D-allopyranose and the X-ray structure of the former
E. Lee, P. Browne, P. McArdle, and D. Cunningham (Galway, Ireland) 285

Regioselective synthesis of α-L-fucosyl-containing disaccharides by use of α-L-fucosidases of various origins
K. Ajisaka and M. Shirakabe (Odawara, Japan) 291

A new route to *C*-glycosyl compounds. Wittig-type reaction promoted by zinc
L. Dheilly, C. Fréchou, D. Beaupère, R. Uzan, and G. Demailly (Amiens, France) 301

The use of chlorodimethylthexylsilane for protecting the hydroxyl groups in cyclomaltoheptaose (β-cyclodextrin)
A. W. Coleman, P. Zhang, C.-C. Ling (Châtenay-Malabry, France), H. Parrot-Lopez, and H. Galons (Paris, France) 307

A facile synthesis of 4-*O*-allyl-D-xylopyranose and its use in the preparation of xylo-oligosaccharides
K. Takeo, Y. Murata, and S. Kitamura (Kyoto, Japan) 311

A new synthesis of 2-deoxy-β-Kdo, a potent CMP-Kdo synthetase inhibitor
H. Ohrui, M. Morita, and H. Meguro (Sendai, Japan) 319

Action of Cu(II) salts on osazones: side reactions of osotriazole formation
H. El Khadem, P. Czubarow, and C. Sandstedt (Washington, DC, U.S.A.) 327

Structural studies of a polysaccharide from the seeds of *Nelumbo nucifera*
S. Das, B. Ray, and P. K. Ghosal (Burdwan, India) 331

Location of the *O*-acetyl group in welan by the reductive-cleavage method
J. D. Stankowski and S. G. Zeller (San Diego, CA, U.S.A.) 337

The effect of the content of D-mannuronic acid and L-guluronic acid blocks in alginates on the antitumor activity
M. Fujihara and T. Nagumo (Kanagawa, Japan) 343

Announcement c1

Author index volume 224 . . . . . c3

Subject index volume 224 . . . . . c5

Contents volume 224 . . . . . c9